



# BIOS RadicalFluid

EC FERTILIZER – COMPOUND FLUID FERTILIZER

N	P	K	U.F.
10	34	0	44

FERTILIZER CE

## PHOSPHORUS AND NITROGEN READILY AVAILABLE

**BIOS RadicalFluid** is a liquid fertilizer specially designed to provide **Nitrogen** and **Phosphorus** to the crops in a perfectly and immediately assimilable form, to stimulate rooting, germination and plant growth.

**BIOS RadicalFluid** is indicated for radical applications as a starter effect and for potassium-based soils.

It can be used by foliar (in case of rescue intervention) to provide highly assimilable phosphorus.

**BIOS RadicalFluid: the right solution for an impeccable start!**

### COMPOSITION

		P/P	equivalent to P/V at 20°C	
Nitrogen	(N) total	10 %	12,5 %	
Nitrogen	(N) ammonia	10 %	12,5 %	
Phosphoric anhydride	(P <sub>2</sub> O <sub>5</sub> ) soluble in water	34 %	42,5 %	



**BIOS RadicalFluid**

### PACKAGING

Cardboard	(Bottle Pz. 20 x Kg.1)
Jug	Kg. 8 - 15 - 30 - 1.300
Form	Fluid

Storage temperature: >= 10° <=30°

Don't expose the container to direct sunlight. The product isn't combustible, and it's stable at ordinary temperatures and pressures. Dispose this product responsibly after use. Avoid contact with skin.

### RAW MATERIALS

### MINERAL COMPOSITION

Fertilizer's solution NP



Restrictions on the use - Art. 11, paragraph 1, letter c) of the Reg.CE/1069/2009: The feeding of farmed animals with herbage, assumed through the pasture or administered after having been picked up, coming from farmland where organic fertilizers or soil improvers different from dung have been applied, unless the pasture or the cut of grass takes after a waiting period - at least 21 days, facing to guarantee a suitable risk assessment for the public and animal health.

### RECOMMENDED DOSES

#### Fertigation

from Kg. 15 to Kg. 60 / Ha

#### Foliar

from Kg. 3 to Kg. 6 for / Ha

Minimum dose level: for repeated use during the cultural cycle

Maximum dose level: effect starter, then the product can be used alone.

The recommended doses have indicative value and should be increased or decreased considering the follow: the pedoclimatic characteristic of the zone of interest, fertility, water retention, structure of soil, cultural variety, the equipment in use and finally the experience of the agricultural entrepreneur. In any case it is recommended to avoid concentrations of the product next to the seed and/or to the roots.



The suitable analytical data on the wrappings follow the prescriptions of the D.L.gs n. 75 of 29/04/2010 and following changes and /or integrations. All the data provided in the present publication are indicative, BIOS s.r.l. the right reserves its rights to modify them without obligation of warning.

FERTILIZER CE – COMPOSED FLUID FERTILIZER